Department of Mathematics
Comprehensive Examination
2023 Spring Semester Core Classes

Time: 2.5 hours
Directions:

-

5. Consider the equation

$$\ln(x) = (x - 2)^2 - 3.$$

(a) Prove that the equation has exactly two solutions.

(b) Let α be the larger of the two solutions and use Newton's Method to find an approximation of α with an absolute error less than 10^{-5}.

6. Consider $A = (a_{ij})$ and its splitting $A = D - L - U$ where

$$D = \text{diag}(a_{11}, a_{22}, \ldots, a_{nn}), \; a_{ii} \neq 0$$

is the diagonal part of A, and

$$L = \begin{bmatrix} 0 & & & \\ a_{21} & 0 & & \\ \vdots & & \ddots & \\ a_{n1} & a_{n2} & \cdots & 0 \end{bmatrix}; \quad U = \begin{bmatrix} 0 & a_{12} & \cdots & a_{1n} \\ & \ddots & & \vdots \\ & & 0 & a_{n-1,n} \\ & & & 0 \end{bmatrix}$$

are the lower and upper triangular parts of A respectively. Consider the following linear system:

$$\begin{aligned} 4x_1 - x_2 - x_3 &= 3 \\ 2x_1 + 6x_2 + x_3 &= 9 \\ x_1 + x_2 + 7x_3 &= 6 \end{aligned}; \quad \mathbf{x}^{(0)} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

(a) For the linear system above, starting with $\mathbf{x}^{(0)}$, find $\mathbf{x}^{(1)}$ and $\mathbf{x}^{(2)}$ using the Jacobi method.

(b) Will the Jacobi method converge for any initial vector?